

6.309 *Euxoa (Euxoa) conspicua* (Hübner, [1827]): Elemento ASM. Según bibliografía univoltina (VI-IX) con diapausa estival. **LF: Ab: 23** (WH56): 23.IX.2006; **25** (WH45): 10.VIII.2005; **125** (XH04): 24.VI.2005. **DV:** 400-1200 m.

6.311 *Euxoa (Euxoa) hastifera* (Donzel, 1847): Elemento ASM. Univoltina (IX). **LF: Ab: 209** (XJ03): 17.IX.2004. **DV:** 640-1480 m.

6.312 *Euxoa (Euxoa) mendelis* Fernández, 1915: Elemento AM. Univoltina (IX). **LF: Ab: 209** (XJ03): 17.IX.2004. **DV:** 640 -1435 m.

6.316 *Euxoa (Euxoa) powelli* (Oberthür, 1912): Elemento AM. En Albacete vuela la ssp. *persubtilis* Corti, 1929. Univoltina (X). **LF: Ab: 25** (WH45): 16.X.2005; **213** (WH41): 2.X.2004. **DV:** 1220-1480 m.

6.318 *Dichagyris candelisequa* (Denis & Schiffermüller, 1775): Elemento EUR. Bivoltina (V y VIII-X). **LF: Ab: 21** (WH55): 1.V.1982, 1.VIII.1983, 31.X.2006; **27** (WH76): 27.VIII.2003; **213** (WH41): 23.VIII.2004, 11.IX.2004. **DV:** 950-1480 m.

6.319 *Dichagyris constanti* (Millière, 1860): Elemento AM. Univoltina (IX). **LF: Ab: 231** (WH51): 21.IX.2003 (A. Albaladejo leg.). **DV:** 1480 m.

6.321 *Dichagyris renigera* (Hübner, [1808]): Elemento ASM. Univoltina (VI-VII). En Albacete vuela la ssp. *murciensis* [Calle, 1983]. **LF: Ab: 17** (WH46): 11.VI.2005; **45** (WH46): 19.VII.2003; **128** (WH51): 30.VI.2001; **213** (WH41): 4.VI.2005. **DV:** 1550-1650 m.

6.326 *Agrotis lasserrei* (Oberthür, 1881): Elemento ASM. Univoltina (X). **LF: Ab: 8** (XH26): 11.X.2006. **DV:** 520 m.

6.331 *Agrotis spinifera* (Hübner, [1808]): Elemento T. En Murcia [9], polivoltina (IV-X). **LF: Ab: 221** (WH95): 31.III.2006. **DV:** 440 m.

6 b) ESPECIES CITADAS EN LA BIBLIOGRAFÍA DE LA PROVINCIA DE ALBACETE

6.2 *Meganola strigula* (Denis & Schiffermüller, 1775): [19]; 6.3 *Meganola togatulalis* (Hübner, 1796): [19]; 6.13 *Eutelia adalatrix* (Hübner, 1813): [6]; 6.14 *Tyta luctuosa* (Denis & Schiffermüller, 1775): [6], [19], [27]; 6.16 *Catephia alchymista* (Denis & Schiffermüller, 1775): [27]; 6.18 *Ophiusa tirhaca* (Cramer, 1777): [6]; 6.19 *Minucia lunaris* (Denis & Schiffermüller, 1775): [3], [4], [6], [19], [27]; 6.20 *Clytie illunaris* (Hübner, [1813]): [6]; 6.21 *Dysgonia algira* (Linnaeus, 1767): [4], [6], [3], [27]; 6.22 *Dysgonia torrida* (Guenée, 1852): [9]; 6.25 *Euclidia glyphica* (Linnaeus, 1758): [3], [4], [6], [27]; 6.26 *Drasteria cailino* (Lefebvre, 1827): [6]; 6.27 *Catocala conjuncta* (Esper, 1787): [4], [20]; 6.28 *Catocala conversa* (Esper, 1787): [17], [4], [19], [20], [27]; 6.29 *Catocala elocata* (Esper, 1787): [4], [6]; 6.30 *Catocala dilecta* (Hübner, [1808]): [6]; 6.32 *Catocala nymphaea* (Esper, 1787): [3], [6], [20]; 6.33 *Catocala nymphagoga* (Esper, 1787): [4], [6], [20]; 6.34